

KRISHZYME™ PROTEIN DEGLYCOSYLATION-NO KIT

REF : KPGF-200

Ver 1.0

RIUO

RIUO

**For Research & Industrial
Use Only**

REF

Catalog Number



Store At

LOT

Batch Code



Manufactured For



Biological Risk



Expiry Date



Consult Operating Instructions

For Research and Industrial Use Only. Purchase does not include or carry the right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of KRISHGEN BioSystems is strictly prohibited.



KRISHGEN BioSystems

Unit Nos#318/319, Shah & Nahar,
Off Dr E Moses Road, Worli, Mumbai 400018.
Tel: 91 (22) 49198700 | Email: sales@krishgen.com

Product Description:


KRISHZYME™ Protein Deglycosylation-NO Kit (for N-linked & Simple O-linked Glycans) contains 1 vial each of PNGase F (Glycerol-free) (Krishzyme™ KPGF-001), O-Glycosidase (Krishzyme™ KPGF-004) and α 2-3,6,8,9Neuraminidase (Krishzyme™ KPGF-005), which can be used simultaneously for the removal of all N-linked Glycans and simple O-linked Glycans (including polysialylated) from glycoproteins.

Krishzyme™ PNGase F (Glycerol-free) cleaves all N-linked complex, hybrid or high mannose oligosaccharides unless α (1,3) core fucosylated (asparagine must be peptide-bonded at both termini).

Krishzyme™ O-Glycosidase, also known as Endo- α -N-Acetylgalactosaminidase, catalyzes the removal of Core 1 and Core 3 O-linked disaccharides from glycoproteins. Unfortunately there is no such “universal” endoglycosidase for O-glycans, these have to be trimmed down to Gal-GalNAc-O-S/T or GlcNAc- β (1-3)-GalNAc-O-S/T before O-glycanase can remove the Core structure. Modification of this core with e.g. sialic acid will block O-glycanase.

Krishzyme™ α 2-3,6,8,9Neuraminidase is included in the kit and can remove α (2,3)-, α (2,6)-, α (2,8)-, and α (2,9)-linked N-acetylneuraminic acid from glycoproteins and oligosaccharides.

One set of this Kit includes 15,000 units of PNGase F (Glycerol-free); 1,200,000 units of O-Glycosidase and 0.6 units of α 2-3,6,8,9 Neuraminidase.



KRISHZYME™ Protein Deglycosylation Kit (for N-linked & Simple O-linked Glycans) can be used simultaneously for the removal of all N-linked Glycans and simple O-linked Glycans (including polysialylated) from glycoproteins.

Product Size :

Cat No	Pack Size	Concentration
KPGF-200	PNGase F(Glycerol-free)	15,000 U / 30 ul
	O-Glycosidase	1,200,000 U / 30 ul
	α 2-3,6,8,9 Neuraminidase	0.6 U / 30 ul

Physical Form:

KRISHZYME™ PNGase F is supplied as a liquid in 20mM Tris-HCl (pH 7.5 at 25°C), 50mM NaCl and 5mM EDTA at a concentration of 500,000 U/ml.

KRISHZYME™ O-Glycosidase is supplied as a liquid in 20mM Tris-HCl (pH 7.5 at 25°C), 50mM NaCl and 1mM EDTA.

KRISHZYME™ α 2-3,6,8,9 Neuraminidase is supplied as a liquid in 20mM Tris-HCl (pH 7.5 at 25°C), 50mM NaCl and 1mM EDTA.

Reagents Supplied:

The following reagents are supplied with this product:

Composition	Formula	Concentration
Denaturing Buffer	5%SDS, 0.4M DTT	10X
Assay Buffer 2	0.5M Sodium Phosphate (pH7.5 at 25°C)	10X
NP-40 Solution		10%

Product Source:

Krishzyme™ PNGase-F (Glycerol free) is cloned from *Elizabethkingia meningoseptica* and expressed in *E.coli*. O-Glycosidase is cloned from *Enterococcus faecalis* and expressed in *E.coli*. α 2-3,6,8,9 Neuraminidase is cloned from *Clostridium perfringens* and overexpressed in *E.coli*.

Product Quality:

≥95% purity, as determined by SDS-PAGE. No other exoglycosidase, endoglycosidase, and protease activity were contaminated.

Unit Definition:

One unit of PNGase F is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 ug of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 ul.

One Unit of O-Glycosidase is defined as the amount of enzyme required to remove 0.68 nmol of O-linked disaccharide from 5mg of neuroaminidase digested, non-denatured fetuin in one hour at 37°C in a total reaction of 100 ul.

One Unit of α 2-3,6,8,9 Neuraminidase is defined as the amount of enzyme required to catalyze the release of 1 umole of p-nitrophenol from p-nitrophenyl- α -D-N-acetylneuroaminic acid per minute at 37°C, pH5.5.

Storage Temperature:

4°C

Characteristic :

- Recombinant enzyme
- Stored in 50% glycerol.
- ≥95% purity, as determined by SDS-PAGE
- Optimal activity and stability for up to 12 months
- Can be used under native or denaturing conditions
- Optimized for deglycosylation of glycoproteins; leaves N-glycan core oligosaccharides intact and suitable for further analysis

Applications:

- Characterizing whether the protein is glycosylated
- Release of intact N-linked glycans from glycopeptides and glycoproteins
- Structure-function studies of N-glycosylated glycoproteins
- Preparation of deglycosylated proteins for molecular weight estimation or crystallography studies

Suggestions for Use:**Denaturing Reaction Conditions:**

- 1) Combine 10-100 ug of glycoprotein, 1 ul of **10X Denaturing Buffer** and **H₂O** (if necessary)

- 2) to make a 30 ul total reaction volume;
- 3) Denature glycoprotein by heating reaction at 100°C for 10 minutes;
- 4) Make a total reaction volume of 50 ul by adding **5 ul of 10XAssay Buffer 2, 5 ul of 10% NP- 40 Solution; 1-2 ul of PNGase-F (glycerol free); 1-2 ul of O-Glycosidase; 1-2 ul of α 2-3,6,8,9 Neuraminidase** and **H₂O (q.s.)**; mix gently;
- 5) Incubate reaction at 37°C for 1-5 hours.
- 6) Use downstream for analysis.

Non-Denaturing Reaction Conditions:

- 1) Combine 10-100 ug of glycoprotein, **5 ul of 10XAssay Buffer 2, 1-2 ul of PNGase-F (glycerol free); 1-2 ul of O-Glycosidase; 1-2 ul of α 2-3,6,8,9 Neuraminidase** and **H₂O (q.s.)** to make total reaction volume of 50 ul; mix gently;
- 2) Incubate reaction at 37°C for 16 hours;
- 3) Use downstream for analysis.

Notes :

- Since KGPF-002 contains 50% glycerol, we recommend limiting PNGase F to 1/10 (or less) of the total reaction volume to keep the final glycerol concentration equal to (or less than) 5%.
- When deglycosylating a native glycoprotein, it is recommended that an aliquot of the glycoprotein is subjected to the denaturing protocol to provide a positive control for the fully deglycosylated protein. The non-denatured reaction can then be compared to the denatured reaction to determine the extent of reaction completion;
- To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required;
- The simplest method of assessing the extent of deglycosylation is by mobility shifts on SDS-PAGE gels;
- Since PNGase F, Recombinant activity is inhibited by SDS, it is essential to have NP- 40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time;
- PNGase F, Recombinant will not cleave N-linked glycans containing core α 1-3 Fucose;
- Recommended Storage Temperature is -20°C.

References:

Characterization of glycoproteins and their associated oligosaccharides through the use of endoglycosidases. ... F Maley, RB Trimble, AL Tarentino... - Analytical biochemistry, 1989 - Elsevier

Deglycosylation of asparagine-linked glycans by peptide: N-glycosidase F. ... AL Tarentino, CM Gomez, TH Plummer Jr - Biochemistry, 1985 - ACS Publications/

Demonstration of peptide: N-glycosidase F activity in endo-beta-N-acetylglucosaminidase F preparations. ... TH Plummer, JH Elder, S Alexander, AW Phelan... - Journal of Biological ..., 1984 - ASBMB

Glycosylation of *Pichia pastoris* -derived proteins. ... RK Bretthauer, FJ Castellino - Biotechnology and applied ..., 1999 - Wiley Online Library.

Pharmacological chaperones rescue cell-surface expression and function of misfolded V2 vasopressin receptor mutants. ... JP Morello, A Salahpour, A Laperrière... - The Journal of ..., 2000 - Am Soc Clin Investig.

A monolithic PNGase F enzyme microreactor enabling glycan mass mapping of glycoproteins by mass spectrometry. ... AK Palm, MV Novotny - ... An International Journal Devoted to the ..., 2005 - Wiley Online Library.

Protein-protein interaction and not glycosylation determines the binding selectivity of heterodimers between the calcitonin receptor-like receptor and the ... S Hilairët, SM Foord, FH Marshall, M Bouvier - Journal of Biological ..., 2001 - ASBMB

Purification and Structure-Function Analysis of Native, PNGase F-Treated, and Endo-. beta.-galactosidase-Treated CHIP28 Water Channels. AN Van Hoek, MC Wiener, JM Verbavatz, D Brown... - Biochemistry, 1995 - ACS Publications

Isolation and characterization of CD47 glycoprotein: a multispanning membrane protein which is the same as integrin-associated protein (IAP) and the ovarian tumour ... WJ Mawby, CH Holmes, DJ Anstee, FA Spring... - Biochemical ..., 1994 - biochemj.org

DC-SIGN and L-SIGN are high affinity binding receptors for hepatitis C virus glycoprotein E2 ..., S Fong, A Amara, C Houles, F Fieschi... - Journal of Biological ..., 2003 - ASBMB

LIMITED WARRANTY

Krishgen Biosystems does not warrant against damages or defects arising in shipping or handling, or out of accident or improper or abnormal use of the Products; against defects in products or components not manufactured by Krishgen Biosystems, or against damages resulting from such non-Krishgen Biosystems made products or components. Krishgen Biosystems passes on to customer the warranty it received (if any) from the maker thereof of such non Krishgen made products or components. This warranty also does not apply to Products to which changes or modifications have been made or attempted by persons other than pursuant to written authorization by Krishgen Biosystems.

THIS WARRANTY IS EXCLUSIVE. The sole and exclusive obligation of Krishgen Biosystems shall be to repair or replace the defective Products in the manner and for the period provided above. Krishgen Biosystems shall not have any other obligation with respect to the Products or any part thereof, whether based on contract, tort, and strict liability or otherwise. Under no circumstances, whether based on this Limited Warranty or otherwise, shall Krishgen Biosystems be liable for incidental, special, or consequential damages.

This Limited Warranty states the entire obligation of Krishgen Biosystems with respect to the Products. If any part of this Limited Warranty is determined to be void or illegal, the remainder shall remain in full force and effect.

Krishgen Biosystems. 2018

THANK YOU FOR USING KRISHGEN PRODUCT!

Disclaimer:

The names of companies indicted and the statements claimed are gleaned from the data sheets and specifications available in open source. They do not in any way reflect an endorsement or a statement by the respective companies.