

Quick Yeast Genomic DNA Extraction Kit

Cat. No: KBN101162

Size: 100 preps

Store at: -20°C

For research use only

For Research Purposes Only. Not for use in diagnostic or therapeutic procedures. Purchase does not include or carry the right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of KRISHGEN BioSystems is strictly prohibited.

KRISHGEN BioSystems Unit Nos#318/319, Shah & Nahar, Off Dr E Moses Road, Worli,
Mumbai 400 018. Tel: (022) 49198700 Email: sales1@krishgen.com

Quick Yeast Genomic DNA Extraction Kit

Description:

The genomic DNA extraction system uses the silica-based membrane technology for simple and fast isolation of genomic DNA (gDNA) without phenol/chloroform. Homogenization is not necessary since tissues are directly lysed by Proteinase K. The buffer system is optimized to allow selective binding of DNA to the silica-based membrane. The simple centrifugation procedure can completely removes impurities such as proteins, divalent cations, and secondary metabolites. Pure DNA is then eluted in water or low-salt buffer, ready to use.

The Kit is suitable for high quality genomic DNA purification from yeast.

Materials Provided:

Solution DS	30 ml
Solution MS	40 ml
Proteinase K (20mg/ml)	2 ml
Wash Buffer PS	60 ml
Wash Buffer PE	30 ml
Elution Buffer TE (10mM Tris-HCl, 1mM EDTA, pH 8.5)	10 ml
Spin Columns	100 each

Downstream Applications:

Purified DNA is free of impurities and enzyme inhibitors, and have an $A_{260/280}=1.7-1.9$, is suitable for applications such as:

- Genotyping
- PCR/qPCR
- Restriction enzyme digestion
- Sequencing
- Southern blotting

Features:

- **Fast** - procedure takes only 60 min
- **High efficiency** - 3-35 ug of genomic DNA from $1-5 \times 10^7$ yeast cells
- **Safe** - no phenol/chloroform extraction step
- **High purity** - purified DNA without enzyme inhibitors, RNA or proteins, ready for downstream applications

Storage:

Store Proteinase K at -20°C, other reagents at room temperature for up to 1 year.

Any precipitate in the Solution DS and Solution MS can be dissolved by incubating at 37°C before use.

Important Notes:

- Prior to the initial use of the kit, dilute the Wash Buffer PE with ethanol (96-100%):

Solution	KBN101162 (100 preps)
Wash Buffer PE	30 ml
Ethanol	90 ml
Total Volume	120 ml

Mix well, mark the labels on the bottle that ethanol is added.

- Ensure that no DNases are introduced into the sterile solutions of the kit.
- Make sure there is no precipitates in Solution DS and Solution MS. If any precipitate is visible, warming the solutions at 37°C for 3-5 min to dissolve the precipitate, and cooling to 25°C before use.
- Wear disposable gloves when handling the Solution MS as it contains guanidine hydrochloride.
- All purification steps should be carried out at room temperature.
- All centrifugations should be carried out by a table-top microcentrifuge at >12,000 g (10,000-14,000 rpm, depending on the rotor type).

Protocol:

1. Add 1-5 ml of yeast cells (not exceed 5×10^7 cells) to a 1.5 ml microcentrifuge tube. Centrifuge at 12,000 rpm for 1 min to pellet the cells. Remove the supernatant.
2. Add 600 ul lyticase buffer and 200U **lyticase**, mix thoroughly by brief vortexing or inverting, incubate at 30°C for 30 min. Centrifuge at 5,000 rpm for 8-10 min. Remove the supernatant.
 - Lyticase need to be supplied by the user
3. Add 200 ul **Solution DS**. Mix immediately and thoroughly by brief vortexing or inverting.
Optional • If RNA-free genomic DNA is required, add 4 ul RNase A (100 mg/ml) and incubate for 5 min at room temperature. RNase A can be purchased separately.
4. Add 20 ul **Proteinase K**, Mix thoroughly by brief vortexing or inverting. Incubate at 55°C until yield a homogeneous solution (~30 min).
5. Add 220 ul **Solution MS**, Mix thoroughly by brief vortexing or inverting. Incubate at 65°C for 10 min (inverting several times to yield a homogeneous solution).
6. Add 220 ul **ethanol (96–100%)** to the lysate, and mix thoroughly by brief vortexing or inverting.
7. Pipet the mixture from step 5 into the spin column placed in a 2 ml collection tube (provided). Centrifuge at 12,000 rpm for 1 min. Discard flow-through.
 - Genomic DNA is adsorbed on the silica membrane of the column in this step.
8. Add 500 ul **Wash Buffer PS**, and centrifuge for 1 min at 12,000 rpm. Discard flow-through.
9. Add 500 ul **Wash Buffer PE**, and centrifuge for 1 min at 12,000 rpm. Discard flow-through.
 - Wash Buffer PE must be diluted with ethanol (96-100%) previously.
10. Repeat step 8.
11. Centrifuge for 3 min at 12,000 rpm to dry the column membrane. Discard flow-through and collection tube.
 - Since residual ethanol may interfere with subsequent reactions, it is important to dry the membrane of the spin column. This centrifugation step ensures that no residual ethanol will be carried during the following elution step. If carryover of ethanol occurs, empty the collection tube, then reuse it after centrifuging for 1 min at 12,000 rpm.
12. Place the spin column in a clean 1.5 ml microcentrifuge tube (not provided), and pipet 30-100 ul **Elution Buffer TE** directly onto the membrane. Incubate at room temperature for 2 min.
 - Elution buffer TE can be replaced by deionized water. But the pH should be 8.0-8.5.
 - Prewarm Elution Buffer TE to 65 °C can increase the yield of genomic DNA.
13. Centrifuge for 2 min at 12,000 rpm. The tube contains the purified DNA. **Store the DNA at -20 °C.**

LIMITED WARRANTY

Krishgen Biosystems does not warrant against damages or defects arising in shipping or handling or out of accident or improper or abnormal use of the products; against defects in products or components not manufactured by Krishgen Biosystems or against damages resulting from such non-Krishgen Biosystems made products or components. Krishgen Biosystems passes on to customer the warranty it received (if any) from the maker of such non Krishgen made products or components. This warranty also does not apply to products to which changes or modifications have been made or attempted by persons other than pursuant to written authorization by Krishgen Biosystems.

THIS WARRANTY IS Exclusive. The sole and exclusive obligation of Krishgen Biosystems shall be to repair or replace the defective products in the manner and for the period provided above. Krishgen Biosystems shall not have any other obligation with respect to the products or any part thereof, whether based on contract, tort, and strict liability or otherwise. Under no circumstances, whether based on this Limited Warranty or otherwise, shall Krishgen Biosystems be liable for incidental, special, or consequential damages. This Limited Warranty states the entire obligation of Krishgen Biosystems with respect to the products. If any part of this Limited Warranty is determined to be void or illegal, the remainder shall remain in full force and effect.

Krishgen Biosystems. 2025

THANK YOU FOR USING A KRISHGEN PRODUCT!